

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/271023042>

# Kinesio taping in musculoskeletal pain and disability that lasts for more than 4 weeks: Is it time to peel off the tape and throw it out with the sweat? A systematic review with me...

Article in *British Journal of Sports Medicine* · January 2015

DOI: 10.1136/bjsports-2014-094151 · Source: PubMed

CITATIONS

54

READS

707

2 authors:



**Edwin Lim**

Singapore Institute of Technology (SIT)

10 PUBLICATIONS 226 CITATIONS

[SEE PROFILE](#)



**Mathew Tay**

La Trobe University

1 PUBLICATION 53 CITATIONS

[SEE PROFILE](#)

# Kinesio taping in musculoskeletal pain and disability that lasts for more than 4 weeks: is it time to peel off the tape and throw it out with the sweat?

## A systematic review with meta-analysis focused on pain and also methods of tape application

Edwin Choon Wyn Lim,<sup>1</sup> Mathew Guo Xiang Tay

<sup>1</sup>Department of Physiotherapy, Singapore General Hospital, Singapore

<sup>2</sup>Collaborative Care (Clinical Services), National Healthcare Group Polyclinics, Singapore

### Correspondence to

Dr Edwin Choon Wyn Lim, Department of Physiotherapy, Singapore General Hospital, Singapore 169608, Singapore; [edwin.lim@uqconnect.edu.au](mailto:edwin.lim@uqconnect.edu.au)

Accepted 26 December 2014

### ABSTRACT

**Introduction** In recent years, Kinesio tape has been used to support injured muscle and joints, and relieve pain. We compared the pain and disability in individuals with chronic musculoskeletal pain who were treated with Kinesio taping with those using minimal or other treatment approaches.

**Methods** Searches of eight major electronic databases were conducted. Data for pain and disability scores were extracted. Meta-analyses (wherever possible) with either a fixed or random effect(s) model, standardised mean differences (SMDs) and tests of heterogeneity were performed.

**Results** Seventeen clinical-controlled trials were identified and included in the meta-analyses. When compared to minimal intervention, Kinesio taping provided superior pain relief (pooled SMD=-0.36, 95% CI -0.64 to -0.09, p=0.009) but the pooled disability scores were not significantly different (pooled SMD=-0.41, 95% CI -0.83 to 0.01, p=0.05). No significant differences were found when comparing Kinesio taping to other treatment approaches for pain (pooled SMD=-0.44, 95% CI -1.69 to 0.82, p=0.49) and disability (pooled SMD=0.08, 95% CI -0.27 to 0.43, p=0.65).

**Discussion** Kinesio taping is superior to minimal intervention for pain relief. Existing evidence does not establish the superiority of Kinesio taping to other treatment approaches to reduce pain and disability for individuals with chronic musculoskeletal pain.

### INTRODUCTION

In recent years, the application of Kinesio tape (KT) has emerged as an interesting and relatively novel method for treating musculoskeletal conditions.<sup>1</sup> This elastic tape purportedly mimics the thickness of the skin and the manufacturers claim it works by lifting the skin, which increases blood circulation and lymphatic drainage leading to a reduction in pain.<sup>2</sup> Ever since it was donated for use during the high-volume physiotherapy treatment at the Olympic Games,<sup>3 4</sup> it is now increasingly seen on high-profile athletes and is being used by healthcare practitioners today.<sup>5</sup>

The bulk of published literature does not favour the use of KT to improve range of motion,<sup>6</sup> strength,<sup>7 8</sup> proprioception<sup>9</sup> and functional performance.<sup>10</sup> However, there have been limited positive findings in terms of range of motion<sup>11 12</sup> and strength.<sup>13</sup> Many systematic reviews<sup>4 14-17</sup> did

not report effectiveness of Kinesio taping for musculoskeletal injuries in clinical practice; however, some reviews<sup>18 19</sup> have reported evidence for pain reduction with the use of KT. From a methodological perspective, many of the reviews<sup>14-16 18</sup> used a descriptive approach to summarise their results, while the others<sup>4 17 19</sup> attempted a quantitative approach to evaluate the effects of KT on certain clinical parameters. Even though there have been extensive efforts invested in evaluating the efficacy of KT, there is still a dearth of attempts in collating the findings of individual studies to determine the effects of KT application on pain and disability, and, if it does, the magnitude of these effects.

We aimed to systematically review randomised controlled trials comparing the effect of KT (intervention) with other forms of interventions (comparisons) for pain and disability (outcomes) in individuals with chronic musculoskeletal pain (participants). We included trials that also had individuals with chronic musculoskeletal pain or musculoskeletal pain persisting beyond the acute phase, defined as pain lasting longer than 4 weeks.<sup>20</sup> A secondary objective was to review the parameters of KT application and to investigate whether these factors influence pain and disability outcomes.

### METHODS

#### Search strategy

We searched MEDLINE (1966-present), CINAHL (1966-present), EMBASE (all years), Cochrane Central Register of Controlled Trials, Physiotherapy Evidence Database (PEDro), Scopus, Google Scholar and ProQuest Dissertations and Theses databases for literature on the use of Kinesio taping for pain and disability in chronic musculoskeletal conditions. The limits applied included 'humans' and 'randomised controlled trials'. The last search was run on 2 July 2014. The following search terms were used to search the databases: kinesio taping; kinesio\* adj2 taping; taping and strapping; and musculoskeletal pain. These steps were then repeated for the other databases. The reviewers followed a selection process, defined prior to the beginning of the review, which included a checklist for inclusion criteria. Articles were eligible for inclusion if they were randomised controlled human trials, included individuals with chronic musculoskeletal pain condition or musculoskeletal

**To cite:** Lim ECW, Tay MGX. *Br J Sports Med* Published Online First: [please include Day Month Year] doi:10.1136/bjsports-2014-094151

pain that persisted beyond the acute phase,<sup>20</sup> assigned an experimental group to receive treatment which includes KT, assigned the comparison group to receive other forms of interventions other than KT and those which used outcome measures that included pain and disability measures. Eligibility assessment for the manuscript was performed by two reviewers (ECWL and MGXT). Disagreements between reviewers were resolved by consensus.

### Data extraction and quality assessment

The methodological quality of the trials was assessed using the 11-item PEDro scale.<sup>21</sup> Assessment of quality of trials was performed by two independent reviewers (ECWL and MGXT). We assessed the methodological quality of the studies by evaluating the domains of population, treatment allocation, blinding, prognostic comparability and analysis. By using a standardised extraction form the information on characteristics of trial participants (age, gender and duration of complaint), details of intervention (amount of tension applied and duration of each tape worn in situ), and pre-outcome and post-outcome measures (pain and disability) was extracted from each included trial. Outcomes that were closer to the end of treatment were used whenever there were multiple time points within a study. If the outcomes were reported for more than one activity, testing condition, or over more than one site, the pre-outcome and post-outcome measures which gave the worst mean difference were extracted. When there was inadequate information about the outcomes to allow data analysis, the authors were contacted. Of the three authors who were contacted, one replied to our queries.<sup>22</sup>

### Quantitative data synthesis and analysis

Reliability analyses of inter-rater agreement were performed with PASW Statistics V.18.0 for Windows (SPSS Inc, Chicago, Illinois, USA). Inter-rater reliability was reported for the total quality score with  $\kappa$  statistics<sup>23</sup> and was interpreted as poor (<0.00), slight (0.00–0.20), fair (0.21–0.40), moderate (0.41–0.60), substantial (0.61–0.80) or almost perfect (0.81–1.0).

Where appropriate and possible, the results were pooled with formal meta-analytical techniques using RevMan 5 (Copenhagen, Denmark: The Nordic Cochrane Centre, The Cochrane Collaboration, 2006). To account for differing outcome scales used in the studies, we calculated the standardised mean differences (SMDs) for pain and disability scores, their 95% CIs and performed tests of heterogeneity ( $\chi^2$ ). The  $I^2$  statistic was used to measure the extent of between-trial heterogeneity. Fixed effects or random effects models were used as appropriate and were based on our interpretation of commonality of effect size.<sup>24</sup> For example, data were pooled using a random effects model, if trials differed in ways that might have plausibly impacted on the pooled outcome.<sup>24</sup>

Subgroup analyses based on intervention (control arm) strata were then analysed to make the control arm more comparable to the arm with KT. For example, trials in which participants received no taping or sham taping when assigned to the control groups were grouped as 'KT versus minimal intervention'. Such grouping also included trials which used other methods of ensuring minimal intervention to the control arm, that is, by providing exercise, general physiotherapy, etc to both groups. In contrast, trials in which participants received treatment (other than KT) when assigned to the control groups were grouped as 'KT versus other forms of intervention'. The differences were calculated such that negative differences indicated that the results favoured KT, while positive differences indicated that the

results favoured the alternate intervention (ie, minimal intervention or other forms of intervention). For all analyses, significance was set at  $p < 0.05$ . To assess the risk of publication bias (resulting from non-publication of small trials with negative results), we also plotted SMD versus SE for both pain and disability scores.<sup>25</sup> The symmetry of this 'funnel plot' was assessed visually.<sup>25</sup> A funnel plot is a scatter plot of intervention effect against a measure of study size. It is used primarily as a visual aid in detecting bias or systematic heterogeneity. A symmetric inverted funnel shape arises from a 'well-behaved' data set in which publication bias is unlikely. An asymmetric funnel indicates a relationship between intervention effect and study size. This suggests the possibility of either publication bias or a systematic difference between smaller and larger studies ('small study effects').

## RESULTS

### Study selection

The initial electronic database search resulted in a total of 606 articles; of these, 29 were selected for detailed inspection and 17 eligible papers remained in this review. [Figure 1](#) shows the flow of papers through review and includes the reasons for exclusion of articles. One additional paper<sup>26</sup> was also discovered during a more detailed reading of these papers and was included.

### Methodological quality


There was substantial agreement between the two reviewers ( $\kappa = 0.62$ ,  $p < 0.001$ ). Percentage of agreement for individual items ranged from 36.4% to 100%. The methodological quality assessment using the PEDro scale revealed a mean score of 6.17 (range 3–9) of a possible 10 points; that is, quality scores for individual studies ranged from 30% to 90% (3–9) of maximum possible attainable points ([table 1](#)). Criteria commonly not met were concealment of allocation, and blinding of treating therapists and patients. It was noted that one of the cross-over randomised trials<sup>27</sup> used insufficient amount of time for 'washout' between treatment periods, which may give rise to a carryover effect<sup>28</sup> and potentially affect the pooled estimates. Consequently, we decided to exclude this paper due to methodological concern.

### Study characteristics

Seventeen trials (416 patients in the experimental group and 406 patients in the control group), which had data for measures of pain and/or disability, were available for pooling ([figure 2](#)). Four trials evaluated the effects of KT on pain and/or disability in knee pain,<sup>29–31 36</sup> four trials in low back pain,<sup>32 34 37 38</sup> three trials in neck pain,<sup>33 35 39</sup> two trials in shoulder pain,<sup>41 42</sup> two trials in plantar fasciitis,<sup>26 40</sup> one trial in de Quervain's disease<sup>22</sup> and one trial in myofascial pain.<sup>43</sup> Evidence of considerable symmetry in the funnel plots ([figure 2A, B](#)) was visually confirmed. The studies were distributed symmetrically about the combined effect size and there was an equal concentration of studies on either side of the SMD.

### Pain—KT versus minimal intervention

Data were pooled using a random effects model; there was a significant pooled SMD in pain ( $-0.68$ , 95% CI  $-1.11$  to  $-0.25$ ,  $p = 0.002$ ) between the KT and minimal intervention groups. There was a high level of heterogeneity ( $I^2 = 80\%$ ,  $\tau^2 = 0.40$ ,  $\chi^2 = 49.39$ ,  $df = 10$ ,  $p < 0.00001$ ; [figure 3A](#)). To investigate whether the presence of outliers could have contributed to the overall significance, we excluded two studies.<sup>30 33</sup> However, the



**Figure 1** Selection process for studies included in analysis.

pooled SMD in pain between the groups remained significant ( $-0.36$ , 95% CI  $-0.64$  to  $-0.09$ ,  $p=0.009$ ) with a low-to-moderate level of heterogeneity ( $I^2=42\%$ ,  $\tau^2=0.07$ ,  $\chi^2=13.87$ ,  $df=8$ ,  $p=0.09$ ).

#### Pain—KT versus other forms of intervention

When comparing SMD in pain between the KT and other intervention groups, the pooled SMD in pain between the groups was not significant ( $-0.44$ , 95% CI  $-1.69$  to  $0.82$ ,  $p=0.49$ ) with a high level of heterogeneity ( $I^2=96\%$ ,  $\tau^2=2.72$ ,  $\chi^2=154.58$ ,  $df=6$ ,  $p<0.00001$ ; [figure 3A](#)).

#### Disability—KT versus minimal intervention

When comparing SMD in disability between the KT and minimal intervention groups, the pooled SMD in disability between the groups was not significant ( $-0.41$ , 95% CI  $-0.83$  to  $0.01$ ,  $p=0.05$ ) with a moderate level of heterogeneity ( $I^2=66\%$ ,  $\tau^2=0.15$ ,  $\chi^2=11.93$ ,  $df=4$ ,  $p=0.02$ ; [figure 3B](#)).

#### Disability—KT versus other forms of intervention

When comparing SMD in disability between the KT and other intervention groups, the pooled SMD in disability between the groups was not significant ( $0.08$ , 95% CI  $-0.27$  to  $0.43$ ,

**Table 1** Details of the included randomised controlled trials

Author (year), study design, population	Intervention (n)	Age (years)	Gender (M, F)	Duration (months), mean (SD)	Brief details of intervention	Preintervention, mean (SD)	Postintervention, mean (SD)	Mean difference between groups (95% CI)	Score
Akbas (2011), <sup>29</sup> RCT, patellofemoral pain syndrome	KT+exercise=15	41.1 (11.3)	M 0, F 15	11.8 (10.8)	Individualised Kinesio taping, 5 strips, changed every 4–5 days for 6 weeks and home exercises (see below)	VAS pain* 2.57 (2.15)	1.71 (1.67)	0.5 (–0.21 to 1.22)	7
	Exercise=16	44.9 (7.75)	M 0, F 16	14.8 (16.3)	Home exercises—6 weeks of stretches, strengthening and open and closed chain exercises	VAS pain* 3.16 (3.98)	0.81 (1.16)		
Anandkumar (2014), <sup>30</sup> RCT, knee osteoarthritis	KT=20	55.7 (5.8)	M 9, F 11	8.4 (1.5)	Kinesio taping, 3 strips, 50–75% tension, origin to insertion, for 30 min	VAS pain 7.4 (1.1)	5.0 (1.3)	–2.07 (–2.85 to –1.28)	7
	Sham=20	55.9 (5.0)	M 8, F 12	8.4 (1.1)	Sham KT, without stretch, for 30 min	VAS pain 7.3 (1.3)	7.5 (1.2)		
Aytar (2011), <sup>31</sup> patellofemoral pain syndrome	KT=12	22.4 (1.62)	M 0, F 12	16.2 (9.68)	Kinesio taping, 4 strips, 50–75% tension, origin to insertion, 45 min	VAS pain* 42.5 (24.5)	42.1 (23.9)	0.14 (–0.7 to 0.98)	6
	Placebo=10	26.2 (3.52)	M 0, F 10	13.7 (8.0)	Placebo—plaster without stretch	VAS pain* 45.0 (23.2)	41.0 (23.8)		
Castro-Sanchez (2012), <sup>32</sup> RCT, low back pain	KT=30	50.0 (15.0)	M 9, F 21	Not reported	Kinesio taping, 4 strips, 25% tension, kept in situ for 1 week	VAS pain 5.6 (1.8) ODI 28.0 (3.0) RMDQ 10.9 (2.1)	4.7 (1.4) 26.0 (3.0) 9.8 (2.2)	VAS pain –0.72 (–1.25 to –0.19) ODI 0.0 (–2.17 to 2.17) RMDQ	8
	Control=29	47.0 (13.0)	M 10, F 19		Sham KT application, 1 strip, kept in situ for a week	VAS pain 5.4 (1.3) ODI 29.0 (3.0) RMDQ 9.8 (2.9)	5.6 (1.4) 27.0 (6.0) 8.6 (3.0)	0.1 (–1.22 to 1.42)	
Dawood (2013), <sup>33</sup> RCT, mechanical neck pain	KT+exercise=19	Range 22–36†	M 39, F 15	At least 3 monthst	Kinesio taping, two strips, replaced every 4 days over 4 weeks and exercises (see below)	VAS pain 7.34 (0.86) NDI 23.87 (9.62)	2.56 (0.75) 9.53 (3.17)	VAS pain KT+exercise vs exercise–2.42 (–3.32 to –1.53) KT+exercise vs cervical traction +exercise	6
	Cervical traction +exercise=19				Cervical traction, 15–20 min, 3 times per week for 4 weeks and exercise (see below)	VAS pain 7.16 (1.20) NDI 22.6 (11.03)	2.86 (0.84) 10.56 (4.5)	KT+exercise vs cervical traction +exercise –0.48 (–1.08 to 0.12) NDI	
	Exercise=16				Exercises—neck stretches, postural correction and isometric exercises	VAS pain 6.98 (0.86) NDI 23.32 (7.57)	4.5 (1.17) 18.6 (7.59)	KT+exercise vs exercise –9.62 (–15.0 to –4.29) KT+exercise vs cervical traction +exercise –2.3 (–8.06 to 3.46)	
Homayouni (2013), <sup>22</sup> RCT, de Quervain's disease	KT=30	45.2 (2.2)	M 13, F 17	2.025 (0.55)	Kinesio taping, 3 strips, stretched by 70%, insertion to origin, repeated four times weekly	VAS pain 58 (6.2)	13 (3.1)	–27.0 (–29.6 to –24.4)	5

Continued

Table 1 Continued

Author (year), study design, population	Intervention (n)	Age (years)	Gender (M, F)	Duration (months), mean (SD)	Brief details of intervention	Preintervention, mean (SD)	Postintervention, mean (SD)	Mean difference between groups (95% CI)	Score
Kachanathu (2014), <sup>34</sup> RCT, low back pain	PT=30	46.0 (1.3)	M 14, F 16	1.825 (0.7)	Paraffin bath, ultrasound, transcutaneous electrical nerve stimulation and friction massage, every 3 days for 10 sessions	VAS pain 56 (5.7)	38 (4.0)		
	Conventional physiotherapy +KT=20	34.8 (7.54)	M 30, F 10	At least 3 months†	Kinesio taping, 2 strips, origin to insertion, three treatment sessions per week for 4 weeks and conventional physiotherapy (see below)	VAS pain 6.2 (1.4) RMDQ 10.3 (3.2)	6.0 (1.8) 10.8 (5.0)	–1.00 (–2.06 to 0.06)	4
Llopis (2012), <sup>35</sup> RCT, mechanical neck pain	Conventional physiotherapy=20				Conventional physiotherapy—stretching exercises, strengthening exercises	VAS pain 2.9 (1.4) RMDQ 4.7 (2.9)	3.7 (2.0) 7.0 (5.5)		
	KT+usual care=5	30.1 (12.0)	Not reported‡	Not reported‡	Kinesio taping, 2 strips, insertion to origin, two treatment sessions per week for 6 weeks and usual care (see below)	Change in VAS pain* 5.55 (1.47)		–0.87 (–2.21 to 0.46)	3
Osorio (2013), <sup>36</sup> cross-over RCT, patellofemoral pain syndrome	Usual care=5				Usual care—stretching exercises, strengthening exercises and massage	Change in VAS pain* 3.98 (1.76)			
	KT=20	21.2 (2.9)	M 7, F 13	Not reported	Kinesio taping over the patella and knee joint, 4 strips, insertion to origin; 2nd and 3rd strips with mild amount of tension	VAS pain* 3.0 (2.2)	1.6 (2.0)	–0.30 (–1.64 to 1.04)	5
Paoloni (2011), <sup>37</sup> RCT, low back pain	McConnell=20				Medial glide McConnell taping	VAS pain* 3.0 (2.2)	1.9 (1.7)		
	KT+Exercise=13	62.0 (13.4)	M 5, F 8	At least 12 weeks†	Kinesio taping, 3 strips, 40% tension, changed every 3 days and therapeutic exercises (see below)	VAS pain 7.6 (1.6) RMDQ 9.5 (4.0)	3.7 (2.5) 7.3 (3.6)	KT vs exercise VAS pain 0.10 (–1.68 to 1.88)	6
	KT=13	62.7 (13.7)	M 5, F 8		Kinesio taping (see above)	VAS pain 7.1 (1.9) RMDQ 10.3 (4.3)	3.1 (2.8) 9.5 (6.8)	RMDQ 3.7 (–0.13 to 7.53)	
Parreira (2014), <sup>38</sup> RCT, low back pain	Exercise=13	62.7 (10.4)	M 4, F 9		Therapeutic exercises—relaxation, and stretching and active exercises	VAS pain 7.6 (1.7) RMDQ 9.9 (3.6)	3.5 (2.4) 5.4 (3.9)		
	KT=74	51.0 (15.0)	M 18, F 56	24.0 (264.0)	Kinesio taping, 2 strips over the erector spinae muscle, 10–15% tension, twice a week, over 4 weeks	VAS Pain 7.0 (2.0) RMDQ 11.5 (6.2)	4.4 (2.8) 8.3 (6.9)	VAS pain –0.17 (–0.49 to 0.16)	9
	Sham=74	50.0 (15.0)	M 15, F 59	36.0 (176.0)	Sham taping, same Kinesio taping but without tension, twice a week, over 4 weeks	VAS Pain 6.8 (2.0) RMDQ 10.4 (5.3)	4.6 (2.5) 7.4 (6.4)	RMDQ –0.2 (–2.22 to 1.82)	
Saavedra-Hernandez (2012), <sup>39</sup> RCT, mechanical neck pain	KT=40	46.0 (9.0)	M 21, F 19	82.0 (19.0)	Kinesio taping, 2 strips, 15–25% stretch, insertion to origin, kept in situ for a week	NPRS 5.2 (1.4) NDI 21.4 (2.3)	2.7 (1.2) 15.4 (1.8)	NPRS –0.20 (–0.91 to 0.51)	9
	Manipulation=36	44.0 (10.0)	M 19, F 17	75.0 (18.0)	Thrust manipulation directed at the mid-cervical spine and cervicothoracic junction	NPRS 5.0 (1.9) NDI	2.7 (1.6)	NDI –0.3 (–1.79 to 1.19)	

Continued

Table 1 Continued


Author (year), study design, population	Intervention (n)	Age (years)	Gender (M, F)	Duration (months), mean (SD)	Brief details of intervention	Preintervention, mean (SD)	Postintervention, mean (SD)	Mean difference between groups (95% CI)	Score
Shahane (2013), <sup>40</sup> RCT, plantar fasciitis	KT+ myofascial release=30	Range 40–60†	M 30, F 30	At least 6 weeks†	Kinesio taping over the ankle and calcaneum, origin to insertion, 5 times per week for 2 weeks. Myofascial release (see below) Myofascial release 15 min per session and plantar fascia stretch, 30 s hold, 6 reps per session. Both were given 5 sessions per week for 2 weeks	VAS pain 7.33 (0.84)	16.8 (3.9)	2.93 (2.46 to 3.4)	5
	Stretches+myofascial release=30					VAS pain 7.30 (0.88)	2.37 (1.00)		
Shakeri (2013), <sup>41</sup> RCT, shoulder impingement syndrome	KT=15	46.5 (13.3)	M 7, F 8	7.63 (7.43)	Kinesio taping, four strips, 50–75% stretch, insertion to origin, replaced every 2–3 days over 1 week	VAS pain* 5.86 (1.8)	2.9 (2.25)	–0.72 (–1.46 to 0.02)	8
	Placebo=15	46.6 (14.2)	M 8, F 7	9.33 (10.5)	Placebo Kinesio taping; three I-strips with no tension	VAS pain* 5.53 (1.55)	4.2 (2.7)		
Simsek (2013), <sup>42</sup> RCT, subacromial impingement syndrome	KT+Ex group=19	48†	M 8, F 11	10.4 (8.26)	Kinesio taping, 3 strips, 50–75% stretch, insertion to origin, changed every 3 days for 12 days, and exercises (rotator cuff strengthening and scapular stabilisation)	VAS Pain* 2.74 (2.73) DASH 46.15 (19.83)	1.43 (2.22)	VAS pain –0.28 (–0.92 to 0.36) DASH –15.4 (–26.8 to, –4.0)	7
	Sham+Ex group=19	53†	M 5, F 14	10.4 (6.65)	Sham taping with exercises (rotator cuff strengthening and scapular stabilisation)	VAS pain* 3.21 (2.92) DASH 52.69 (16.42)	2.65 (2.67)		
Tsai (2010), <sup>26</sup> RCT, plantar fasciitis	KT+PT group=29	52.7 (28.8)	M 19, F 33	3.92 (1.80)	Kinesio taping, 2 strips, 33% stretch, insertion to origin, kept in situ for a week, and physical therapy program (see below)	MMPQ 9.29 (2.69)	4.14 (3.02)	–0.72 (–1.26 to –0.19)	5
	PT group=28	30.5 (13.1)		4.33 (3.01)	Therapeutic ultrasound, low-frequency electrotherapy	MMPQ 14.63 (2.61)	11.88 (2.36)		
van der Westhuizen (2012), <sup>43</sup> RCT, myofascial pain	KT=25	27.1 (6.20)	M 7, F 18	Not reported	Kinesio taping, anchor and three halved strips, 15–25% stretch, insertion to origin, replaced every 3 days over 1 week	NRS pain 36.7 (17.8) NDI 8.04 (3.62)	11.8 (13.6)	NRS pain –9.92 (–19.6 to –0.22) NDI 0.88 (–1.43 to 3.19)	6
	Dry needling=25	28.3 (6.16)	M 9, F 16		Dry needling over upper trapezius	NRS pain 32.9 (19.7) NDI 8.96 (4.3)	3.88 (4.55) 18.0 (17.7) 3.92 (4.03)		

\*Values which gave worst outcome (mean difference) were extracted; when pain measures were reported in more than 1 activity, condition, testing condition or site.

†Mean and/or SD not reported.

‡Gender not reported.

DASH, Disabilities of the Arm, Shoulder and Hand (score out of 100); F, female; KT, Kinesio tape; M, male; MMPQ, McGill Melnick pain questionnaire (score out of 20); NDI, Neck Disability Index (score out of 50), with higher scores indicating greater disability; NPIS, Numeric Pain Intensity Scale (score out of 10); NPRS, Numerical pain rating scale (score out of 10); ODI, Oswestry Disability Index (score out of 100); RCT, randomised controlled trial; RMDQ, Roland-Morris Disability Questionnaire (score out of 24); SPADI, Shoulder Pain and Disability Index (score out of 100), with higher scores indicating greater pain and disability; VAS, visual analogue scale (score out of 10 cm or 100 mm).



**Figure 2** Funnel plot of standardised mean difference (SMD) against SE based on (A) pain intensity and (B) disability.

$p=0.65$ ) with a low level of heterogeneity ( $I^2=29\%$ ,  $\tau^2=0.04$ ,  $\chi^2=4.23$ ,  $df=3$ ,  $p=0.24$ ; [figure 3B](#)).

### Parameters of KT application

Four studies (23.5%)<sup>30 31 41 42</sup> used an average tension of 62.5% stretch and four studies (23.5%)<sup>35 37 42 43</sup> replaced the tape every 3 days ([table 2](#)). The average amount of tension applied ranged from 12.5% to 70% stretch,<sup>22 26 29–32 37–39 41–43</sup> while the average duration of each tape worn in situ ranged from 45 min to 7 days<sup>22 26 29 31–33 35 37–43</sup> ([table 2](#)).

Multivariable meta-regression analysis identified amount of tension applied (% stretch) ( $\beta=0.796$ ,  $p=0.024$ ) and duration

of each tape applied in situ (days) ( $\beta=0.644$ ,  $p=0.046$ ) as independent predictors of the effect size of the reported pain score in participants with chronic musculoskeletal pain who received KT ([table 3](#)). The model correctly predicted 80.7% of the effect size of reported pain score. Owing to the limited number of studies which measured disability, multivariable meta-regression analyses could not be computed.

### DISCUSSION


This systematic review synthesised the evidence for KT effectiveness in participants with painful conditions who are managed in primary healthcare practices. KT proved to be superior to minimal intervention, which includes no taping, sham taping and usual care, for the reduction of pain in individuals with >4 weeks of musculoskeletal pain. However, KT is not more effective than other forms of intervention in reducing pain. KT is also not more effective than minimal or other forms of intervention in the reduction of disability related to chronic musculoskeletal pain.

Besides reflecting the multidimensional phenomenon of disability, which encompasses the complex interaction between impairment, activity limitation and participation restriction,<sup>44</sup> the null effect of KT on disability signals the need for conventional therapy, for example, other active forms of intervention, over and above passive treatment, in the management of musculoskeletal pain lasting >4 weeks. This is supported by findings of non-significant differences in pain and/or disability between groups reported by trials which used Kinesio taping only.<sup>32 38</sup>

### KT as an adjunct to exercise therapy

In contrast, the two trials which used KT as an adjunct to exercise reported a significant difference in pain and disability between groups, favouring the experimental group.<sup>33 42</sup> This is not surprising as exercise has been reported to garner good evidence of effectiveness as a standalone or adjunctive treatment for chronic musculoskeletal pain.<sup>45</sup> Taken together, our review suggests that KT, when used in combination with conventional therapy, may be effective in reducing pain. Our clinical impression is that many clinicians use KT in this way—as an adjunct to exercise.

Compared to minimal intervention, we found a significant effect size (reduction in pain) by 0.36 to 0.68, which is considered small to moderate, among patients with >4 weeks of



**Figure 3** Forest plot (effect size and 95% CI) of the (A) pain and (B) disability score of randomised controlled trials. Pooled estimates which represent effect size (standardised mean difference) and 95% CI (using a random effects model) are indicated by empty symbols (KT, Kinesio tape).



**Table 2** The parameters of Kinesio taping used by the various studies (n=17)

Parameters	Significant difference in outcome measure (favouring Kinesio taping)			
	Pain		Disability	
	Yes	No	Yes	No
Average amount of tension used (% stretch)				
12.5		Low back pain <sup>38</sup>		Low back pain <sup>38</sup>
20		Myofascial pain; <sup>43</sup> mechanical neck pain <sup>39</sup>		Mechanical neck pain, <sup>39</sup> myofascial pain <sup>43</sup>
25	Low back pain <sup>32</sup>			Low back pain <sup>32</sup>
33	Plantar fasciitis <sup>26</sup>			
35		Patellofemoral pain syndrome <sup>29</sup>		
40		Low back pain <sup>37</sup>		Low back pain <sup>37</sup>
62.5	Knee osteoarthritis <sup>30</sup>	Patellofemoral pain syndrome; <sup>31</sup> shoulder pain <sup>41 42</sup>	Shoulder pain <sup>42</sup>	
70	de Quervain's disease <sup>22</sup>			
Not reported	Plantar fasciitis; <sup>40</sup> mechanical neck pain <sup>33</sup>	Mechanical neck pain; <sup>35</sup> patellofemoral pain syndrome <sup>36</sup> Low back pain <sup>34</sup>	Mechanical neck pain <sup>33</sup>	Low back pain <sup>34</sup>
Average duration of each tape worn in situ (days)				
<2	Knee osteoarthritis <sup>30</sup> Plantar fasciitis <sup>40</sup> de Quervain's disease <sup>22</sup>	Patellofemoral pain syndrome <sup>31</sup>		
2		Low back pain <sup>38</sup>		Low back pain <sup>38</sup>
2.5		Shoulder pain <sup>41</sup>		
3		Shoulder pain, <sup>42</sup> mechanical neck pain, <sup>35</sup> myofascial pain <sup>43</sup> Low back pain <sup>37</sup>	Shoulder pain <sup>42</sup>	Low back pain, <sup>37</sup> myofascial pain <sup>43</sup>
4	Mechanical neck pain <sup>33</sup>		Mechanical neck pain <sup>33</sup>	
5		Patellofemoral pain syndrome <sup>29</sup>		
7	Plantar fasciitis; <sup>26</sup> low back pain <sup>32</sup>	Mechanical neck pain <sup>39</sup>		Low back pain; <sup>32</sup> mechanical neck pain <sup>39</sup>
Not reported		Patellofemoral pain syndrome <sup>36</sup> Low back pain <sup>34</sup>		Low back pain <sup>34</sup>

musculoskeletal pain who received KT. We propose that this observed magnitude is a clinically meaningful change.<sup>46</sup> This is borne out to some extent by the mean difference of 0.92 to 1.33 during an ad hoc analysis of the weighted mean difference in pain measure between groups.

Our finding did not concur with a recent systematic review<sup>19</sup> which reported a non-significant negligible magnitude of 0.08. The difference in results could be attributed to the inclusion of trials which were non-randomised<sup>47 48</sup> and trials which sampled participants with acute musculoskeletal conditions<sup>12 49</sup> in their meta-analysis.<sup>19</sup> For example, individuals with varying duration of symptoms may respond differently to KT, such that the effects in different directions for the acute pain and more long-lasting pain subgroups may combine to give the impression of no effect for the combined group (washout effect).<sup>50</sup>

### What is the mechanism of action?

To date, the underlying pain-relief mechanism of such proprietary tape remains poorly understood. Nonetheless, it is tempting to speculate that the afferent stimuli provided by the applied KT may inhibit the transmission of nociceptive signals to the spinal level<sup>51</sup> in chronic musculoskeletal pain conditions, leading to the attenuation of pain experience. The basis of such assertion is the commonality in feature, that is, spinal cord hyperexcitability among chronic musculoskeletal pain conditions.<sup>52</sup> However, the approximate thickness of the KT in relation to the epidermis of the skin was intended to avoid sensory stimuli when properly applied.<sup>53</sup> To this end, it remains unclear

to what extent the gate control theory is involved in the efficacy of KT.

On another note, it is plausible that the decongestive property of KT may play a role, at least in part, in the efficacy of KT on pain reduction among musculoskeletal pain lasting >4 weeks with swelling. Such assertion is inferred from two of the trials with large effect sizes,<sup>22 30</sup> that is, Anandkumar *et al*<sup>30</sup> who investigated the efficacy of KT in knee osteoarthritis (OA) which is characterised by pain and joint effusion,<sup>54</sup> while Homayouni *et al*<sup>22</sup> investigated chronic de Quervain's disease which is characterised by tenderness and swelling around the styloid process.<sup>55</sup> A possible explanation for these observations could be the skin-lifting effect of KT, which promotes blood circulation and improves lymph drainage<sup>53</sup> leading to a reduction in swelling and subsequently, to pain-relief. Conversely, Stedge *et al*<sup>56</sup> reported a lack of significant difference in the blood circulation of the gastrocnemius muscle between their KT group, and sham or no tape groups in a healthy active population.

It is also conceivable that the applied KT reduces pain by stimulating the descending pain inhibitory mechanism from the higher centres of the brain.<sup>19</sup> This is corroborated, at least in part, by the significant reduction in pain observed in three<sup>30 32 42</sup> of six placebo-controlled studies.<sup>30–32 38 41 42</sup> Other proposed mechanisms include the lifting of the skin by the applied KT, resulting in a reduction of pressure on the subcutaneous nociceptors.<sup>57</sup> Apart from the aforementioned biophysiological explanations, it is plausible that KT may exert its effect via the bioneuro-psychological pathway.<sup>58</sup> Recently, Montalvo *et al*<sup>19</sup> drew attention to the probable

**Table 3** Predictors for Kinesio taping treatment effect size in chronic musculoskeletal pain conditions

Dependent variables	Independent variables	Unstandardised coefficients		95% CI for B		Standardised coefficients		t Value	Significance
		B	SE	Lower	Upper	$\beta$			
Standardised mean difference of reported pain score	Amount of tension applied (% stretch)	0.038	0.011	0.008	0.067	0.796		3.524	0.024
	Duration of each tape applied in situ (days)	0.255	0.089	0.007	0.503	0.644		2.853	0.046
	Constant	-3.577	0.563	-5.139	-2.014			-6.357	0.003

95% CI, 95% CI for regression coefficient;  $\beta$ ,  $\beta$  coefficient; B, regression coefficient; SE, SE of B; t, t statistics.

release of analgesic neurotransmitters due to the placebo effect, that is, patient's expectations.<sup>59–61</sup> It remains uncertain to what degree the KT serves as a psychological crutch.<sup>62</sup> Further research is warranted to confirm these postulated mechanisms.

### Application of KT

Our review of the literature revealed variability in KT application and that it is uncertain if the tape can be applied with a greater pain-relieving effect. Importantly, five studies<sup>33–36 40</sup> did not report the amount of tension applied while two<sup>34 36</sup> did not report the duration of each tape worn in situ. Notwithstanding, our review suggests that the effect size (SMD) for pain reduction was lower when the studies applied more tension and left the applied tape in situ longer.

Our finding extends that of Kase *et al*<sup>53</sup> who specified that excessive tension may diminish the beneficial effects of KT.<sup>53</sup> In addition, they mentioned that the longitudinal elasticity of the KT is effective for 3–5 days,<sup>53</sup> implying that the applied tape should be replaced, at the most, every 5 days. However, these parameters were not identified as predictors of effect size (SMD) for pain reduction in ad hoc univariate meta-regression analyses. Moreover, the results are limited to associations rather than causations between the variables, and this may act as a caveat on inferences drawn from our findings.

Limitations of the review include the small number of studies and relatively small sample size used in the multivariable meta-regression analysis. Also, our results are drawn from studies with somewhat varying methodology and treatment 'dosage' of KT application. For example, eight studies<sup>22 26 35 36 39 41–43</sup> applied the KT from the insertion to the origin of the muscle, four studies<sup>30 31 34 40</sup> taped it from the origin to insertion direction, while the remaining studies<sup>29 32 33 37 38</sup> did not report details regarding the taping direction. Such variation may have influenced, at least in part, the outcome measures since a muscle facilitatory or inhibitory effect would be elicited depending on the taping direction.<sup>53</sup> Furthermore, the individuals with musculoskeletal pain >4 weeks in the included studies received a single to 12 treatments sessions<sup>22 26 30–43</sup> of KT at a frequency of one to five times per week,<sup>22 26 29 32–35 37–43</sup> over a span of 1–6 weeks.<sup>22 26 29 32–35 37–43</sup> There is a need for the use of standard application procedures (by body parts) to determine the efficacy of KT in future studies.

Future studies may consider recruiting patients who respond to KT prior to randomisation. On another methodological note, it is worth mentioning that none of the included trials measured the patient's expectations or beliefs regarding the benefits of Kinesio taping with the use of simple validated questionnaires. This would be pertinent to therapy studies as it has been reported to greatly influence the outcomes among those with

musculoskeletal pain.<sup>58 63</sup> Publication bias might account for the effect size we observed in this review. However, a symmetrical funnel plot suggests that missing studies or even selective reporting within included studies are unlikely. Moreover, we did not apply language restriction during our search. The high level of clinical heterogeneity among the trials, with respect to the different clinical conditions, poses a challenge during this review. In addition, musculoskeletal pain may not be sufficiently representative in this review because knee OA was represented by only one paper<sup>30</sup> and many other musculoskeletal pain states have not been evaluated. Therefore, variations in KT application across the studies may limit overall conclusions.

In conclusion, this review highlights that KT is superior to minimal intervention for pain relief. Existing evidence does not establish the superiority of KT in reducing disability when compared to either minimal or other forms of intervention. Taken together, our review indicates that KT, when used in combination with conventional therapy, may be effective in reducing pain.

### What are the new findings

- ▶ Kinesio tape (KT) is superior to minimal intervention for pain relief.
- ▶ Our evidence does not establish superiority of KT over other treatment approaches in reducing pain and disability for individuals with musculoskeletal pain >4 weeks.
- ▶ The amount of tension applied and duration of applied tape left in situ may influence the effect size for pain reduction.

**Contributors** ECWL contributed to conception, design, construct of the study and the analyses. ECWL and MGXT conducted the comprehensive search and critical appraisal and contributed to the manuscript writing. ECWL is the guarantor.

**Competing interests** None.

**Provenance and peer review** Not commissioned; externally peer reviewed.

### REFERENCES

- 1 Kamper SJ, Henschke N. Kinesio taping for sports injuries. *Br J Sports Med* 2013;47:1128–9.
- 2 Kinesio Taping® Association International: Kinesio taping® method. 2013. <http://www.kinesiotaping.com/about/kinesio-taping-method> (accessed 21 Dec 2014).
- 3 Grant ME, Steffen K, Glasgow P, *et al*. The role of sports physiotherapy at the London 2012 Olympic Games. *Br J Sports Med* 2014;48:63–70.
- 4 Williams S, Whatman C, Hume PA, *et al*. Kinesio taping in treatment and prevention of sports injuries: a meta-analysis of the evidence for its effectiveness. *Sports Med* 2012;42:153–64.
- 5 Kinesio UK short-listed for Sports technology Awards. 2014. <http://www.kinesiotaping.co.uk/news/articles/2014-02-28-Kinesio-UK-short-listed-for-Sports-Technology-Awards/index.jsp> (accessed 6 Aug 2014).

## Review

- 6 Gomez-Soriano J, Abian-Vicen J, Aparicio-Garcia C, *et al*. The effects of Kinesio taping on muscle tone in healthy subjects: a double-blind, placebo-controlled crossover trial. *Man Ther* 2014;19:131–6.
- 7 Csapo R, Alegre LM. Effects of Kinesio taping on skeletal muscle strength—a meta-analysis of current evidence. *J Sci Med Sport* Published Online First: 27 Jun 2014.
- 8 Chang HY, Cheng SC, Lin CC, *et al*. The effectiveness of Kinesio taping for athletes with medial elbow epicondylar tendinopathy. *Int J Sports Med* 2013;34:1003–6.
- 9 Halseth T, McChesney JW, Debeliso M, *et al*. The effects of Kinesio taping on proprioception at the ankle. *J Sports Sci Med* 2004;3:1–7.
- 10 Schiffer T, Mollinger A, Sperlich B, *et al*. Kinesio taping and jump performance in female elite track and field athletes. *J Sport Rehabil* Published Online First: 23 Jun 2014.
- 11 Yoshida A, Kahanov L. The effect of Kinesio taping on lower trunk range of motions. *Res Sports Med* 2007;15:103–12.
- 12 Gonzalez-Iglesias J, Fernandez-de-Las-Penas C, Cleland JA, *et al*. Short-term effects of cervical Kinesio taping on pain and cervical range of motion in patients with acute whiplash injury: a randomized clinical trial. *J Orthop Sports Phys Ther* 2009;39:515–21.
- 13 Hsu YH, Chen WY, Lin HC, *et al*. The effects of taping on scapular kinematics and muscle performance in baseball players with shoulder impingement syndrome. *J Electromyogr Kinesiol* 2009;19:1092–9.
- 14 Bassett K, Lingman S, Ellis R. The use and treatment efficacy of kinaesthetic taping for musculoskeletal conditions: a systematic review. *N Z J Physiother* 2010;38:56.
- 15 Mostafavifar M, Wertz J, Borchers J. A systematic review of the effectiveness of Kinesio taping for musculoskeletal injury. *Phys Sportsmed* 2012;40:33–40.
- 16 Morris D, Jones D, Ryan H, *et al*. The clinical effects of Kinesio(R) Tex taping: a systematic review. *Physiother Theory Pract* 2013;29:259–70.
- 17 Parreira Pdo C, Costa Lda C, Hespagnol Junior LC, *et al*. Current evidence does not support the use of Kinesio taping in clinical practice: a systematic review. *J Physiother* 2014;60:31–9.
- 18 Kairon A, Bar-Sela S. A systematic review of the effectiveness of Kinesio Taping—fact or fashion? *Eur J Phys Rehabil Med* 2013;49:699–709.
- 19 Montalvo AM, Cara EL, Myer GD. Effect of kinesiology taping on pain in individuals with musculoskeletal injuries: systematic review and meta-analysis. *Phys Sportsmed* 2014;42:48–57.
- 20 Chou R, Huffman LH. Medications for acute and chronic low back pain: a review of the evidence for an American Pain Society/American College of Physicians clinical practice guideline. *Ann Intern Med* 2007;147:505–14.
- 21 de Morton NA. The PEDro scale is a valid measure of the methodological quality of clinical trials: a demographic study. *Aust J Physiother* 2009;55:129–33.
- 22 Homayouni K, Zeynali L, Mianehsaz E. Comparison between Kinesio taping and physiotherapy in the treatment of de Quervain's disease. *J Musculoskelet Res* 2013;16:1–6.
- 23 Landis JR, Koch GG. The measurement of observer agreement for categorical data. *Biometrics* 1977;33:159–74.
- 24 Borenstein M, Hedges L, Higgins J, *et al*. *Introduction to meta-analysis*. Chichester, West Sussex, UK: Wiley, 2009.
- 25 Liberati A, Altman DG, Tetzlaff J, *et al*. The PRISMA statement for reporting systematic reviews and meta-analyses of studies that evaluate health care interventions: explanation and elaboration. *Ann Intern Med* 2009;151:W65–94.
- 26 Tsai CT, Chang WD, Lee JP. Effects of short-term treatment with kinesiotaping for plantar fasciitis. *J Musculoskelet Pain* 2010;18:71–80.
- 27 Campolo M, Babu J, Dmochowska K, *et al*. A comparison of two taping techniques (Kinesio and McConnell) and their effect on anterior knee pain during functional activities. *Int J Sports Phys Ther* 2013;8:105–10.
- 28 Elbourne DR, Altman DG, Higgins JP, *et al*. Meta-analyses involving cross-over trials: methodological issues. *Int J Epidemiol* 2002;31:140–9.
- 29 Akbas E, Atay AO, Yuksel I. The effects of additional Kinesio taping over exercise in the treatment of patellofemoral pain syndrome. *Acta Orthop Traumatol Turc* 2011;45:335–41.
- 30 Anandkumar S, Sudarshan S, Nagpal P. Efficacy of Kinesio taping on isokinetic quadriceps torque in knee osteoarthritis: a double blinded randomized controlled study. *Physiother Theory Pract* 2014;30:375–83.
- 31 Aytaç A, Ozunlu N, Surenkok O, *et al*. Initial effects of Kinesio® taping in patients with patellofemoral pain syndrome: a randomized, double-blind study. *Isokinet Exerc Sci* 2011;19:135–42.
- 32 Castro-Sanchez AM, Lara-Palomo IC, Mataran-Penarrocha GA, *et al*. Kinesio taping reduces disability and pain slightly in chronic non-specific low back pain: a randomised trial. *J Physiother* 2012;58:89–95.
- 33 Dawood RS, Kattabei OM, Nasef SA, *et al*. Effectiveness of Kinesio taping versus cervical traction on mechanical neck dysfunction. *Int J Ther Rehabil Res* 2013;2:1–5.
- 34 Kachanathu SJ, Alenazi AM, Seif HE, *et al*. Comparison between Kinesio taping and a Traditional Physical Therapy Program in treatment of nonspecific low back pain. *J Phys Ther Sci* 2014;26:1185–8.
- 35 Llopis LG, Aranda MC. Intervención fisioterápica con vendaje neuromuscular en pacientes con cervicalgia mecánica. Un estudio piloto. *Fisioterapia* 2012;34:189–95.
- 36 Osorio JA, Vairo GL, Rozea GD, *et al*. The effects of two therapeutic patellofemoral taping techniques on strength, endurance, and pain responses. *Phys Ther Sport* 2013;14:199–206.
- 37 Paoloni M, Bernetti A, Fratocchi G, *et al*. Kinesio taping applied to lumbar muscles influences clinical and electromyographic characteristics in chronic low back pain patients. *Eur J Phys Rehabil Med* 2011;47:237–44.
- 38 Parreira Pdo C, Costa Lda C, Takahashi R, *et al*. Kinesio Taping to generate skin convolutions is not better than sham taping for people with chronic non-specific low back pain: a randomised trial. *J Physiother* 2014;60:90–6.
- 39 Saavedra-Hernandez M, Castro-Sanchez AM, Arroyo-Morales M, *et al*. Short-term effects of Kinesio taping versus cervical thrust manipulation in patients with mechanical neck pain: a randomized clinical trial. *J Orthop Sports Phys Ther* 2012;42:724–30.
- 40 Shahane S. *The effect of myofascial release technique and stretching versus myofascial release technique and taping in patients with chronic plantar fasciitis: a comparative study* [thesis]. Bangalore, Karnataka, India: Rajiv Gandhi University of Health Sciences, 2013.
- 41 Shakeri H, Keshavarz R, Arab AM, *et al*. Clinical effectiveness of kinesiological taping on pain and pain-free shoulder range of motion in patients with shoulder impingement syndrome: a randomized, double blinded, placebo-controlled trial. *Int J Sports Phys Ther* 2013;3:800.
- 42 Simsek HH, Balki S, Keklik SS, *et al*. Does Kinesio taping in addition to exercise therapy improve the outcomes in subacromial impingement syndrome? A randomized, double-blind, controlled clinical trial. *Acta Orthop Traumatol Turc* 2013;47:104–10.
- 43 van der Westhuizen JH. *The relative effectiveness of Kinesiotape® versus dry needling in patients with myofascial pain syndrome of the Trapezius muscle* [thesis]. Durban University of Technology, 2014.
- 44 Disabilities. 2014. <http://www.who.int/topics/disabilities/en/> (accessed 14 Sep 2014).
- 45 Hassett AL, Williams DA. Non-pharmacological treatment of chronic widespread musculoskeletal pain. *Best Pract Res Clin Rheumatol* 2011; 25:299–309.
- 46 Salaffi F, Stancati A, Silvestri CA, *et al*. Minimal clinically important changes in chronic musculoskeletal pain intensity measured on a numerical rating scale. *Eur J Pain* 2004;8:283–91.
- 47 Kaya E, Zinnuroglu M, Tugcu I. Kinesio taping compared to physical therapy modalities for the treatment of shoulder impingement syndrome. *Clin Rheumatol* 2011;30:201–7.
- 48 Kuru T, Yaliman A, Dereli EE. Comparison of efficiency of Kinesio(R) taping and electrical stimulation in patients with patellofemoral pain syndrome. *Acta Orthop Traumatol Turc* 2012;46:385–92.
- 49 Thelen MD, Dauber JA, Stoneman PD. The clinical efficacy of Kinesio tape for shoulder pain: a randomized, double-blinded, clinical trial. *J Orthop Sports Phys Ther* 2008;38:389–95.
- 50 Straker LM, O'Sullivan PB, Smith AJ, *et al*. Relationships between prolonged neck/shoulder pain and sitting spinal posture in male and female adolescents. *Man Ther* 2009;14:321–9.
- 51 DeLeo JA. Basic science of pain. *J Bone Joint Surg Am* 2006;88(Suppl 2):58–62.
- 52 Lim ECW, Sterling M, Stone A, *et al*. Central hyperexcitability as measured with nociceptive flexor reflex threshold in chronic musculoskeletal pain: a systematic review. *Pain* 2011;152:1811–20.
- 53 Kase K, Wallis J, Kase T. *Clinical therapeutic applications of the Kinesio taping method*. Tokyo, Japan: Ken Ikai, 2003.
- 54 Hunter DJ, McDougall JJ, Keefe FJ. The symptoms of osteoarthritis and the genesis of pain. *Med Clin North Am* 2009;93:83–100, xi.
- 55 Kay NR. De Quervain's disease. Changing pathology or changing perception? *J Hand Surg Br* 2000;25:65–9.
- 56 Stedje HL, Kroskie RM, Docherty CL. Kinesio taping and the circulation and endurance ratio of the gastrocnemius muscle. *J Athl Train* 2012;47:635–42.
- 57 Kahanov L. Kinesio taping (R), part 1: an overview of its use in athletes. *Athl Ther Today* 2007;12:17–18.
- 58 Bialosky JE, Bishop MD, Cleland JA. Individual expectation: an overlooked, but pertinent, factor in the treatment of individuals experiencing musculoskeletal pain. *Phys Ther* 2010;90:1345–55.
- 59 Kamper SJ, Williams CM. The placebo effect: powerful, powerless or redundant? *Br J Sports Med* 2013;47:6–9.
- 60 Oken BS. Placebo effects: clinical aspects and neurobiology. *Brain* 2008;131:2812–23.
- 61 Bishop MD, Mintken PE, Bialosky JE, *et al*. Patient expectations of benefit from interventions for neck pain and resulting influence on outcomes. *J Orthop Sports Phys Ther* 2013;43:457–65.
- 62 Vercelli S, Ferriero G, Bravini E, *et al*. How much is Kinesio taping a psychological crutch? *Man Ther* 2013;18:e11.
- 63 Vercelli S, Sartorio F, Foti C, *et al*. Immediate effects of kinesiotaping on quadriceps muscle strength: a single-blind, placebo-controlled crossover trial. *Clin J Sport Med* 2012;22:319–26.



# Kinesio taping in musculoskeletal pain and disability that lasts for more than 4 weeks: is it time to peel off the tape and throw it out with the sweat? A systematic review with meta-analysis focused on pain and also methods of tape application

Edwin Choon Wyn Lim and Mathew Guo Xiang Tay

*Br J Sports Med* published online January 16, 2015

---

Updated information and services can be found at:  
<http://bjsm.bmj.com/content/early/2015/01/16/bjsports-2014-094151>

---

## References

*These include:*

This article cites 53 articles, 8 of which you can access for free at:  
<http://bjsm.bmj.com/content/early/2015/01/16/bjsports-2014-094151#BIBL>

## Email alerting service

Receive free email alerts when new articles cite this article. Sign up in the box at the top right corner of the online article.

---

## Notes

---

To request permissions go to:  
<http://group.bmj.com/group/rights-licensing/permissions>

To order reprints go to:  
<http://journals.bmj.com/cgi/reprintform>

To subscribe to BMJ go to:  
<http://group.bmj.com/subscribe/>